Generalized Riemann Problems in Computational Fluid Dynamics

MATANIA BEN-ARTZI and JOSEPH FALCOVITZ

The Hebrew University of Jerusalem



PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS

The Edinburgh Building, Cambridge CB2 2RU, UK 40 West 20th Street, New York, NY 10011-4211, USA 477 Williamstown Road, Port Melbourne, VIC 3207, Australia Ruiz de Alarcón 13, 28014 Madrid, Spain Dock House, The Waterfront, Cape Town 8001, South Africa

http://www.cambridge.org

© Cambridge University Press 2003

This book is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2003

Printed in the United Kingdom at the University Press, Cambridge

Typeface Times Roman 10/13 pt. System LATEX $2_{\mathcal{E}}$ [TB]

A catalog record for this book is available from the British Library.

Library of Congress Cataloging in Publication Data

Ben-Artzi, Matania, 1948-

Generalized Riemann problems in computational fluid dynamics / Matania Ben-Artzi, Joseph Falcovitz.

p. cm. – (Cambridge monographs on applied and computational mathematics; 11)

Includes bibliographical references and index.

ISBN 0-521-77296-6

Fluid dynamics. 2. Riemann-Hilbert problems. I. Falcovitz, Joseph, 1937–.
 II. Title. III. Series.

QA911 .B36 2003

532'.05 - dc21 2002034811

ISBN 0521772966 hardback

Contents

| List | ist of Figures | | page xi |
|------|----------------|---|---------|
| Pre | face | - | XV |
| 1 | Intro | oduction | 1 |
| I | BAS | IC THEORY | 5 |
| 2 | Scala | ar Conservation Laws | 7 |
| | 2.1 | Theoretical Background | 7 |
| | 2.2 | Basic Concepts of Numerical Approximation | 25 |
| 3 | The | GRP Method for Scalar Conservation Laws | 36 |
| | 3.1 | From Godunov to the GRP Method | 36 |
| | 3.2 | 1-D Sample Problems | 49 |
| | | 3.2.1 The Linear Conservation Law | 49 |
| | | 3.2.2 The Burgers Nonlinear Conservation Law | 55 |
| | 3.3 | 2-D Sample Problems | 63 |
| 4 | Syste | ems of Conservation Laws | 81 |
| | 4.1 | Nonlinear Hyperbolic Systems in One Space Dimension | 81 |
| | 4.2 | Euler Equations of Quasi-1-D, Compressible, | |
| | | Inviscid Flow | 101 |
| 5 | The | Generalized Riemann Problem (GRP) for | |
| | Com | pressible Fluid Dynamics | 135 |
| | 5.1 | The GRP for Quasi-1-D, Compressible, Inviscid Flow | 135 |
| | 5.2 | The GRP Numerical Method for Quasi-1-D, | |
| | | Compressible Inviscid Flow | 169 |

viii Contents

| 6 | Ana | llytical and Numerical Treatment of Fluid | |
|----|-----|--|-----|
| | Dyn | namical Problems | 184 |
| | 6.1 | The Shock Tube Problem | 184 |
| | 6.2 | Wave Interactions | 189 |
| | | 6.2.1 Shock–Contact Interaction | 192 |
| | | 6.2.2 Shock–Shock Interaction | 195 |
| | | 6.2.3 Shock–CRW Interaction | 203 |
| | | 6.2.4 CRW–Contact Interaction | 207 |
| | 6.3 | Spherically Converging Flow of Cold Gas | 218 |
| | 6.4 | The Flow Induced by an Expanding Sphere | 219 |
| | 6.5 | Converging–Diverging Nozzle Flow | 222 |
| II | NU | MERICAL IMPLEMENTATION | 233 |
| 7 | Fro | m the GRP Algorithm to Scientific Computing | 235 |
| | 7.1 | General Discussion | 235 |
| | 7.2 | Strang's Operator-Splitting Method | 237 |
| | 7.3 | Two-Dimensional Flow in Cartesian Coordinates | 244 |
| 8 | | metric Extensions | 251 |
| | 8.1 | Grids That Move in Time | 251 |
| | 8.2 | e ; e | 252 |
| | 8.3 | £ , £ , , | 255 |
| | | 8.3.1 Basic Setup | 257 |
| | | 8.3.2 Survey of the Full MBT Algorithm | 264 |
| | | 8.3.3 An Example: Shock Lifting of an | |
| | | Elliptic Disk | 266 |
| 9 | A P | hysical Extension: Reacting Flow | 269 |
| | 9.1 | The Equations of Compressible Reacting Flow | 271 |
| | 9.2 | The Chapman–Jouguet (C–J) Model | 276 |
| | 9.3 | The Z–N–D (Zeldovich–von Neumann–Döring) | |
| | | Solution | 281 |
| | 9.4 | The Linear GRP for the Reacting-Flow System | 286 |
| | 9.5 | The GRP Scheme for Reacting Flow | 298 |
| 10 | Way | ve Interaction in a Duct – A Comparative Study | 305 |
| A | Ent | ropy Conditions for Scalar Conservation Laws | 313 |
| В | Con | vergence of the Godunov Scheme | 320 |

| | Contents | ix |
|----------|--|-----|
| C | Riemann Solver for a γ -Law Gas | 330 |
| D | The MUSCL Scheme | 333 |
| Bib | bliography | 337 |
| Glossary | | 345 |
| Index | | 347 |

List of Figures

| 2.1 | Characteristic curves in the nonlinear case. | page 10 |
|------|---|---------|
| 2.2 | A jump discontinuity of a weak solution. | 15 |
| 2.3 | Characteristics for initial step function in Burgers' equation. | 18 |
| 2.4 | Rarefaction wave solution for Burgers' equation. | 19 |
| 2.5 | A jump discontinuity satisfying the entropy condition. | 22 |
| 2.6 | Backward-difference characteristic scheme. | 33 |
| 3.1 | Wave pattern for piecewise-constant initial data. | 38 |
| 3.2 | Wave pattern for the GRP algorithm. | 43 |
| 3.3 | First GRP time-integration cycle of a moving step. | 47 |
| 3.4 | The "slope limiter" is a five-point rule. | 48 |
| 3.5 | First-order integration of $u_t + u_x = 0$, with $u_0 = \sin^4(\pi x)$. | 50 |
| 3.6 | First-order integration of $u_t + u_x = 0$, with step-function | |
| | initial data. | 51 |
| 3.7 | Second-order integration of $u_t + u_x = 0$, with $u_0(x) = \sin^4(\pi x)$ | . 53 |
| 3.8 | Second-order integration of $u_t + u_x = 0$, with step-function | |
| | initial data. | 54 |
| 3.9 | First-order integration of $u_t + (\frac{1}{2}u^2)_x = 0$, with | |
| | $u_0(x) = \sin(2\pi x).$ | 57 |
| 3.10 | First-order integration of $u_t + (\frac{1}{2}u^2)_x = 0$, with step-function | |
| | initial data. | 59 |
| 3.11 | Second-order integration of $u_t + (\frac{1}{2}u^2)_x = 0$, with | |
| | $u_0(x) = \sin(2\pi x).$ | 62 |
| 3.12 | Second-order integration of $u_t + (\frac{1}{2}u^2)_x = 0$, with step-function | |
| | initial data. | 63 |
| 3.13 | Linear diagonal step – Godunov solution. | 67 |
| 3.14 | Linear diagonal step – GRP solution. | 68 |
| 3.15 | Nonlinear diagonal shock – Godunov solution. | 70 |
| | | |

| 3.16 | Nonlinear diagonal shock – GRP solution. | 71 |
|------|--|-----|
| 3.17 | Exact solutions at $t = 1$ of 2-D Burgers equation. | 72 |
| 3.18 | <i>U</i> -level curves for the 2-D Burgers equation at $t = 1$ for Case A. | 73 |
| 3.19 | <i>U</i> -level curves for the 2-D Burgers equation at $t = 1$ for Case B. | 74 |
| 3.20 | <i>U</i> -level curves for the 2-D Burgers equation at $t = 1$ for Case C. | 75 |
| 3.21 | Numerical solutions of 2-D Burgers equation at $t = 1$ for Case D. | 76 |
| 3.22 | The Guckenheimer structure for $u_t + (u^2/2)_x + (u^3/3)_y = 0$. | 77 |
| 3.23 | U-level curves for Guckenheimer equation at $t = 1$. | 79 |
| 4.1 | A simple wave. | 84 |
| 4.2 | A centered rarefaction wave. | 85 |
| 4.3 | States connected to $\tilde{\mathbf{u}}$ by a <i>k</i> -CRW. | 87 |
| 4.4 | States connected to $\tilde{\mathbf{u}}$ by a jump discontinuity. | 91 |
| 4.5 | Shock admissibility conditions. | 94 |
| 4.6 | A contact discontinuity. | 96 |
| 4.7 | The solution to the Riemann problem. | 99 |
| 4.8 | Neighborhood of state \mathbf{u}_R "covered" by interaction curves. | 100 |
| 4.9 | Quasi-1-D flow in a duct. | 102 |
| 4.10 | Right-facing (σ_3) and left-facing (σ_1) shocks. | 113 |
| 4.11 | The thermodynamic Hugoniot curve P_3^+ . | 119 |
| 4.12 | The 3-Hugoniot curve in the (u, p) plane. | 121 |
| 4.13 | The 1-Hugoniot curve in the (u, p) plane. | 121 |
| 4.14 | A planar 1-CRW. It is necessarily isentropic. | 123 |
| 4.15 | A 1-CRW in Lagrangian representation. | 125 |
| 4.16 | A schematic solution to the RP for planar flow. | 127 |
| 4.17 | $u-p$ representation of I_1^l , I_3^r interaction curves. | 129 |
| 4.18 | Solutions to the Riemann problem. | 130 |
| 5.1 | The solutions to the GRP (a), and its associated RP (b). | 137 |
| 5.2 | Structure of the solution to the GRP in Lagrangian coordinates. | 139 |
| 5.3 | The "acoustic case" $V_L = V_R, V_L' \neq V_R'$. | 143 |
| 5.4 | Characteristic coordinates in a 1-CRW. | 146 |
| 5.5 | Possible Eulerian GRP solutions for the wave pattern | |
| | of Figure 5.2. | 163 |
| 5.6 | Structure of the solution for Godunov's scheme. | 172 |
| 5.7 | Distribution of flow variables at time $t = t_n$ (GRP setup). | 173 |
| 6.1 | Eulerian solution of shock tube flow at $t = 20$ (100 cells). | 186 |
| 6.2 | Lagrangian solution of shock tube flow at $t = 20$ (100 cells). | 187 |
| 6.3 | ALE solution of shock tube flow at $t = 20$ (100 cells). | 188 |
| 6.4 | Wave interaction – A schematic description. | 190 |
| 6.5 | Schematic (x, t) diagram of the interaction of two waves. | 190 |
| 6.6 | Riemann problem for the shock–contact interaction. | 193 |

| | List of Figures | xiii |
|------|--|------|
| 6.7 | Wave diagram of the shock–contact interaction. | 193 |
| 6.8 | Initial data for shock—contact interaction. | 194 |
| 6.9 | Shock-contact interaction. Shock and contact merge | |
| | at $(x, t) = (0, 0)$. | 196 |
| 6.10 | Shock—contact postinteraction results at $t = 20$. | 197 |
| 6.11 | Riemann problem for the two-shock interaction. | 198 |
| 6.12 | Wave diagram of the two-shock interaction. | 199 |
| 6.13 | Initial data for shock–shock interaction. | 200 |
| 6.14 | Shock–shock interaction. Shocks merge at $(x, t) = (0, 0)$. | 201 |
| 6.15 | Shock–shock postinteraction results at $t = 18$. | 202 |
| 6.16 | Diagram of the interaction of a 3-shock and a 1-CRW. | 203 |
| 6.17 | Riemann problem for shock–rarefaction interaction. | 204 |
| 6.18 | Initial data for shock–CRW interaction. | 205 |
| 6.19 | Shock–CRW results at preinteraction time $t = -2$. | 206 |
| 6.20 | Shock–CRW postinteraction results at $t = 10$. | 208 |
| 6.21 | Shock–CRW interaction. Time sequence of density. | 209 |
| 6.22 | Interaction of CRW with a contact discontinuity. | 210 |
| 6.23 | CRW-contact interaction schematic. | 210 |
| 6.24 | CRW-contact interaction. Initial data. | 212 |
| 6.25 | CRW-contact interaction. Asymptotic RP initial data. | 214 |
| 6.26 | CRW–contact interaction. $t = 10$. | 215 |
| 6.27 | CRW–contact interaction. $t = 20$. | 216 |
| 6.28 | CRW-contact interaction. Time sequence of density. | 217 |
| 6.29 | Distribution of flow variables for Noh's problem. | 220 |
| 6.30 | Flow in fluid surrounding an expanding sphere. | 223 |
| 6.31 | Laval nozzle flow. Case (A): Godunov scheme. | 226 |
| 6.32 | Laval nozzle flow. Case (A): GRP scheme. | 227 |
| 6.33 | Laval nozzle flow. Case (B): Godunov scheme. | 228 |
| 6.34 | Laval nozzle flow. Case (B): GRP scheme. | 229 |
| 7.1 | Setup for the Euler split scheme. | 248 |
| 8.1 | Boundary polygons and underlying Cartesian grid. | 257 |
| 8.2 | Cell (j, k) intersected by a moving boundary. | 258 |
| 8.3 | Cell (j, k) intersected by a boundary moving in the x direction. | 261 |
| 8.4 | Boundary moving one time step in the <i>x</i> direction. | 265 |
| 8.5 | Kickoff of an elliptic disk by an $M_s = 3$ shock in air. | 267 |
| 9.1 | The C–J Hugoniot curve $H^{(0)}(\tau, p) = 0$. | 278 |
| 9.2 | The Z–N–D solution. | 282 |
| 9.3 | The Z–N–D profile. | 285 |
| 9.4 | GRP computation of ozone decomposition wave. | |
| | $\Delta x = 5 \times 10^{-6}.$ | 302 |

| 9.5 | GRP computation of ozone decomposition wave. | |
|------|---|-----|
| | $\Delta x = 2 \times 10^{-5}.$ | 302 |
| 9.6 | GRP computation of ozone decomposition wave. | |
| | $\Delta x = 5 \times 10^{-5}.$ | 303 |
| 9.7 | Weak detonation wave of ozone decomposition. | 303 |
| 10.1 | CRW interaction with a converging segment. | 308 |
| 10.2 | Comparison of 2-D and quasi-1-D solutions. | 309 |
| B.1 | The piecewise-constant function $U_k(x, t)$. | 323 |
| C.1 | Shock/rarefaction type determination on interaction curve $I_3^{\rm r}$. | 331 |
| C.2 | Solution of RP by intersection between I_1^1 and I_2^r . | 332 |

Introduction

This monograph deals with the *generalized Riemann problem* (GRP) of mathematical fluid dynamics and its application to computational fluid dynamics. It shows how the solution to this problem serves as a basic tool in the construction of a robust numerical scheme that can be successfully implemented in a wide variety of fluid dynamical topics. The flows covered by this exposition may be quite different in nature, yet they share some common features; they all belong to the class of compressible, inviscid, time-dependent flows. Fluid dynamical phenomena of this type often contain a number of smooth flow regions separated by singularities such as shock fronts, detonation waves, interfaces, and centered rarefaction waves. One must then address various computational issues related to this class of fluid dynamical problems, notably the "capturing" of discontinuities such as shock fronts, detonation waves, or interfaces; resolution of centered rarefaction waves where flow gradients are unbounded; and evaluation of flow variables in irregular computational cells at the intersection of a moving boundary surface with an underlying mesh.

From the mathematical point of view, the various systems of equations governing compressible, inviscid, time-dependent flow phenomena may all be characterized as systems of "(nonlinear) hyperbolic conservation laws."

Hyperbolic conservation laws (in one space variable) are systems of time-dependent partial differential equations. The most common problem associated with such systems is the *initial value problem* (henceforth IVP), which is the following: Given the values of the unknown functions at time t=0 (as functions of the space variable $x \in \mathbb{R}$), use the equations to determine the evolution in time of those functions. When the unknown functions are defined over the whole real line \mathbb{R} , one often refers to the IVP as the "Cauchy Problem." In contrast, when the unknown functions are defined only over a finite interval $\mathcal{D} \subseteq \mathbb{R}$, suitable "boundary conditions" must be imposed at the endpoints of \mathcal{D} . From

the physical point of view the latter is clearly the more realistic case. Thus, for example, \mathcal{D} can represent a pipe of finite length, in which one studies the evolution (in time) of flow variables subject to the system of fluid dynamical equations. In this case, the boundary conditions consist of influx and outflux requirements imposed on the pressure, velocity, etc. at the edges of the pipe.

The solutions to the problems considered here possess one common fundamental property, that of "finite propagation speed"; that is, the waves travel at finite speeds. Mathematically speaking, when a change in the initial data is confined to the neighborhood of some point A, it is "felt" by the solution at any other point B only after a certain amount of time, an amount that depends on the distance between the points. It is precisely this feature that allows the construction of "conservation law schemes" for the (numerical) approximation of the solutions.

Although this monograph focuses on the resolution of compressible, inviscid flow problems, and the construction of suitable conservation law schemes, an effort is made to place the treatment in the broader (theoretical and numerical) perspective of hyperbolic conservation laws. However, the necessary background material from physics is also included. We refer the reader to the classical book by R. Courant and K. O. Friedrichs [30] for a thorough discussion of the mathematical aspects of compressible flow. This book also discusses in detail the derivation of the flow equations from the underlying physical conservation laws. For mathematical treatments of hyperbolic conservation laws, we refer to the books by Courant and Hilbert [31], Evans [36], Hörmander [63], Lax [75], and Smoller [103].

To simplify the discussion, we consider primarily the associated Cauchy problem, thus avoiding the further mathematical complications introduced by boundary conditions. Naturally, when dealing with real flow examples, boundary conditions will be needed, and the ways in which they are introduced into the numerical scheme will be explained.

The origin of the subject matter of this monograph can be traced back some forty years, to the early days of computational fluid dynamics. It can best be described by the opening sentence to Section 12.15 of the book by Richtmyer and Morton [96]: "In 1959, Godunov described an ingenious method for one-dimensional problems with shocks."

Godunov's method, as much as it was recognized for its novelty and robustness, suffered from some significant drawbacks. It was Bram van Leer, some twenty years later, who, in an important breakthrough [112], has shown how to modify Godunov's original construction and, indeed, has made it possible to implement the method as the most efficient tool (to date) in this area of computational fluid dynamics. In the simpler case of a scalar conservation law, these ideas will be explained in Chapter 3.

The monograph is divided into two parts, Part I (Basic Theory) and Part II (Numerical Implementation). Part I (Chapters 2–6) deals with the more basic aspects (theoretical and numerical) of systems of conservation laws and the development of the GRP method. Part II (Chapters 7–10) is devoted to several extensions (physical and geometric) of the GRP method for computational fluid dynamics. A more detailed discussion of the contents will follow. The reader will also find a brief summary at the beginning of each chapter.

In writing this monograph we have aimed at a wide readership, consisting not only of graduate students and researchers in applied mathematics but also of those working in various areas of physics and engineering. Yet, we have attempted to maintain a solid level of mathematical rigor. Notions such as "weak solutions" and "convergence of a scheme" are carefully introduced (Chapter 2) in suitable functional settings. We believe that, given the current mathematical level of modern numerical analysis, such concepts ought to be familiar to anyone working in this field. In particular, theorems related to the convergence of the Godunov scheme (in the scalar case) are proved in all mathematical detail (Section 2.2 and Appendix B). In this context we introduce (and compare numerically) some of the "classical" discrete schemes of hyperbolic equations, such as the Lax-Friedrichs and the Lax-Wendroff schemes. At the same time, our main objective in Chapters 2 and 3 is the introduction of the "high-resolution GRP scheme," by way of the Riemann and generalized Riemann problems. We refer to LeVeque [81] for introductory material on finite-difference schemes for conservation laws and to Richtmyer and Morton [96] for the general theory of finite-difference methods (primarily linear theory). A comprehensive survey of the convergence properties of finite-difference schemes to scalar conservation laws can be found in Godlewski and Raviart [54].

In Chapter 4 we introduce systems of conservation laws. The first section outlines the general mathematical background and can be skipped on first reading, as it is of a more mathematical nature. The physical systems of interest, those representing the basic conservation laws of compressible, inviscid flow in the "quasi-one-dimensional" setting, are introduced in the second section. This section is self-contained; the analysis of centered rarefaction waves, as well as the Rankine–Hugoniot shock conditions and the solution to the Riemann problem, is discussed in detail.

Chapter 5 is devoted to the analysis of the GRP in the context of the systems considered in Section 4.2. In Section 5.1 we study the solution to the linear GRP, which is the core of the GRP method. Given linear initial distributions

of the flow variables on the two sides of a jump discontinuity, we determine their instantaneous time derivatives (at that singularity). Van Leer's idea to use this solution for the refinement of Godunov's scheme is implemented in the development of the GRP scheme in Section 5.2.

Chapter 6 is devoted to an investigation of the GRP scheme for fluid dynamics. Numerical results are compared to analytical or asymptotic solutions for a variety of wave interaction problems.

In Chapter 7 we introduce, in rather general terms, the operator-splitting method of Strang. It enables us to extend the GRP algorithm to two-dimensional (2-D) settings, while retaining its second-order accuracy. Chapter 8 deals with further geometric extensions, such as (one-dimensional) "tracking" of singularities and (2-D) moving boundaries.

In Chapter 9 we consider a reacting flow system. The basic set of conservation laws is augmented by a chemical reaction-rate equation, thus providing a simple model of combustion. The GRP algorithm is applied to this extended system.

As a concluding (numerical) example for this monograph, we consider in Chapter 10 a case of wave interaction with a segment of decreasing cross-sectional area in a two-dimensional duct. The major (GRP) numerical approaches developed in Chapters 5, 7, and 8, namely the quasi-1-D approximation and the fully 2-D scheme, are applied to this case. A comparative study of the two solutions sheds light on the nature of the fluid dynamical interaction, as well as on the nature of the quasi-1-D approximation.

Finally, a comment about the numbering system in this book. For the reader's convenience, all theorems, remarks, definitions, claims, etc. within each chapter are sequentially numbered. Thus, for example, Remark 2.22 comes after Definition 2.21 and is followed by Example 2.23.